



**Product Data Sheet** 

# **BIOAKSXTER® M32 Photovoltaic Greenhouse**

#### **Effects**

- Repowering of plant programming capacity.
- Increase in crop yield and reduction in waste production.
- Equilibrium of the vegetation, constant and homogeneous production.
- Increase in vegetative luxuriance and luster of the leaves.
- Increase in the plant's selfdefenses against fungal, viral, bacterial and other diseases.
- Overcoming climate shock and stress conditions.
- Better oxygen exchange and lymphatic flow, with consequential cellular regeneration and tissue elasticity.
- Purification of plant and soil from the pesticides residues.

#### **Product**

BioAksxter® M32 Photovoltaic Greenhouse is the magnetic flux fertilizer, designed for the strengthening of the light uptake program. Thanks to its advanced technology that reproduces natural processes, it programs the equilibrium of plant and soil, and it regenerates vegetal structures by increasing plant potential. In addition, BioAksxter® depollutes and revitalizes the soil, substantially increasing the quality and quantity of yields.

It acts especially on vegetative development, and by strengthening self-defenses it prevents and reduces plant diseases.

It is ideal for best results on leafy vegetable crops, floriculture and nurseries of young plants in all situations of light deficiency.

#### **Action**

BioAksxter® M32 Photovoltaic Greenhouse reproduces the processes of natural fertilization that occurs through water-soil-plant-air exchanges.

BioAksxter® M32 Photovoltaic Greenhouse is an inorganic compound that, when diluted in water, forms a pre-organic compound: it revitalizes soil through self-reproduction of "virgin" microorganisms, and it is readily assimilable by the plant of which it favors "hydrophony" of nutrients, plant's programming capacity, oxygen exchanges, light absorption, regulation of lymphatic flows, and finally cell regeneration.

All this translates into greater plant and soil vitality and ultimately in the best productive results

## **Unique Depolluting Technology**


BioAksxter® M32 Photovoltaic Greenhouse makes up for the insufficient low-impact supply chain because the simple introduction of this advanced technology into cultivation practices closes the circuit of input contaminant molecules, bringing crops and soil back into balance. Its use eliminates chemical residues in agricultural products, it lowers basic radioactivity by preventing alterations in the plant's primary program, and consequently, phenomena of deformation and loss of self-defenses. In the soil, pollutants are transformed into organic absorption structures, and it is possible to notice a rebalancing in the microbiological activity as biodegraders are triggered. BioAksxter® M32 Photovoltaic Greenhouse is not limited to low environmental impact, it goes further, opening the frontier of depollution.

## **Energy for Light Capture**

In crops grown in photovoltaic greenhouses and in all situations of light deficiency, BioAksxter® M32 Photovoltaic Greenhouse reinforces the uptake capacity of plants by accelerating all their photosynthetic processes.

It compensates for up to 70% of diffused light deficiencies, allowing productive results equal to those obtained with direct light.

It allows crops to defend themselves from electromagnetic interference generated by photovoltaic panels. BioAksxter® M32 Photovoltaic Greenhouse enables the increase in yields and the resistance below the minimum threshold of optimal temperatures, prolonging the harvest cycle and slowing the natural deterioration of crops



### Technical Information BIOAKSXTER® M32 Photovoltaic Greenhouse

Capture	accelerating all their photosynthetic processes.  In photovoltaic greenhouses and in all situations of light deficiency, BioAksxter® allows the increase in yields and the resistance below the minimum threshold of the optimal temperatures, prolonging the harvesting cycle and slowing down the natural deterioration of the crops.						
		nce below the minimum threshold o	of the optimal temperatures, prolonging the				
		nce below the minimum threshold o	of the optimal temperatures, prolonging the				
	yields and the resistance below the minimum threshold of the optimal temperatures, prolonging the						
	VIELUS AND THE LESISTAL						
		slowing down the natural deteriorati	on of the crops.				
		9					
He to a	harvesting cycle and s		A 1.*				
How to use	harvesting cycle and s		Application				
How to use	harvesting cycle and s		Application				
Harrida maa	harvesting cycle and s		A				
		9					
		9					
		stowing down the natural deteriorati	on of the crops.				
		slowing down the natural deteriorati	on of the crops.				
	VIELUS AND THE TESISTAL						
	vields and the resistar						
	In photovoltaic greenhouses and in all situations of light deficiency, BioAksxter® allows the increase in yields and the resistance below the minimum threshold of the optimal temperatures, prolonging the						
				Capture			
				Capture			
				Capture			
Capture							
Capture							
Capture							
Capture							
Capture							
Capture							
Captaro							
					In photovoltaic greenhouses and in all situations of light deficiency, BioAksxter® allows the increase in		
					yields and the resistance below the minimum threshold of the optimal temperatures, prolonging the		
					slowing down the natural deteriorati	on of the crops.	
					stowing down the natural detendrati	on on the crops.	
						·	
How to use	harvesting cycle and s		Application				
How to use		Dose per hectare	Application				
How to use	harvesting cycle and s  Type of crop	Dose per hectare					
How to use	Type of crop Floriculture	Dose per hectare 1.5 liters	Application Every 10 days				
How to use	Type of crop Floriculture	Dose per hectare 1.5 liters					
How to use	Type of crop Floriculture Nurseries of young p	Dose per hectare 1.5 liters	Every 10 days				
How to use	Type of crop Floriculture	Dose per hectare 1.5 liters					
How to use	Type of crop Floriculture Nurseries of young p	Dose per hectare 1.5 liters	Every 10 days				
How to use	Type of crop Floriculture Nurseries of young p Leafy greens	Dose per hectare  1.5 liters  1.5 liters	Every 10 days  Every 10 days				
How to use	Type of crop Floriculture Nurseries of young p Leafy greens	Dose per hectare  1.5 liters  1.5 liters	Every 10 days  Every 10 days				
How to use	Type of crop Floriculture Nurseries of young p Leafy greens	Dose per hectare  1.5 liters  1.5 liters	Every 10 days  Every 10 days				
How to use	Type of crop Floriculture Nurseries of young p Leafy greens	Dose per hectare  1.5 liters  1.5 liters	Every 10 days  Every 10 days				
How to use	Type of crop Floriculture Nurseries of young p Leafy greens	Dose per hectare  1.5 liters  1.5 liters	Every 10 days				
How to use	Type of crop Floriculture Nurseries of young p Leafy greens The number of interven	Dose per hectare  1.5 liters  1.5 liters  1.5 liters entions and dosage are indicative ar	Every 10 days  Every 10 days				
How to use	Type of crop Floriculture Nurseries of young p Leafy greens The number of interven	Dose per hectare  1.5 liters  1.5 liters  1.5 liters entions and dosage are indicative ar	Every 10 days  Every 10 days				
How to use	Type of crop Floriculture Nurseries of young p Leafy greens The number of interven	Dose per hectare  1.5 liters  1.5 liters  1.5 liters entions and dosage are indicative ar	Every 10 days  Every 10 days				
How to use	Type of crop Floriculture Nurseries of young p Leafy greens	Dose per hectare  1.5 liters  1.5 liters  1.5 liters entions and dosage are indicative ar	Every 10 days  Every 10 days				
How to use	Type of crop Floriculture Nurseries of young p Leafy greens The number of interventies context, type of cr	Dose per hectare  1.5 liters  1.5 liters  2.5 liters entions and dosage are indicative arop and its duration.	Every 10 days  Every 10 days				
How to use	Type of crop Floriculture Nurseries of young p Leafy greens The number of interventies context, type of cr	Dose per hectare  1.5 liters  1.5 liters  2.5 liters entions and dosage are indicative arop and its duration.	Every 10 days  Every 10 days				
How to use	Type of crop Floriculture Nurseries of young p Leafy greens The number of interventies context, type of cr	Dose per hectare  1.5 liters  1.5 liters  1.5 liters entions and dosage are indicative ar	Every 10 days  Every 10 days				
How to use	Type of crop Floriculture Nurseries of young p Leafy greens The number of interve the context, type of cr Doses can be diluted	Dose per hectare  1.5 liters  1.5 liters  2.5 liters  entions and dosage are indicative are op and its duration. in any volume of water.	Every 10 days  Every 10 days  nd vary, according to instructions, based on				
How to use	Type of crop Floriculture Nurseries of young p Leafy greens The number of interve the context, type of cr Doses can be diluted	Dose per hectare  1.5 liters  1.5 liters  2.5 liters  entions and dosage are indicative are op and its duration. in any volume of water.	Every 10 days  Every 10 days  and vary, according to instructions, based on				
How to use	Type of crop Floriculture Nurseries of young p Leafy greens The number of interve the context, type of cr Doses can be diluted	Dose per hectare  1.5 liters  1.5 liters  2.5 liters  entions and dosage are indicative are op and its duration. in any volume of water.	Every 10 days  Every 10 days  and vary, according to instructions, based on				
How to use	Type of crop Floriculture Nurseries of young p Leafy greens The number of interve the context, type of cr Doses can be diluted In the case of crops a	Dose per hectare  1.5 liters  1.5 liters  entions and dosage are indicative ar op and its duration. in any volume of water. ffected by resistant plant diseases, if	Every 10 days  Every 10 days  d vary, according to instructions, based on  t is essential to double the dosage of the fire				
How to use	Type of crop Floriculture Nurseries of young p Leafy greens The number of interve the context, type of cr Doses can be diluted In the case of crops a	Dose per hectare  1.5 liters  1.5 liters  entions and dosage are indicative ar op and its duration. in any volume of water. ffected by resistant plant diseases, if	Every 10 days  Every 10 days  d vary, according to instructions, based on  t is essential to double the dosage of the fire				
How to use	Type of crop Floriculture Nurseries of young p Leafy greens The number of interve the context, type of cr Doses can be diluted In the case of crops a	Dose per hectare  1.5 liters  1.5 liters  entions and dosage are indicative ar op and its duration. in any volume of water. ffected by resistant plant diseases, if	Every 10 days  Every 10 days  d vary, according to instructions, based on				
How to use	Type of crop Floriculture Nurseries of young p Leafy greens The number of intervethe context, type of croposes can be diluted In the case of crops a 3 interventions. In additional context and some context are context.	Dose per hectare  1.5 liters  1.5 liters  2.5 liters  entions and dosage are indicative ar op and its duration. in any volume of water. ffected by resistant plant diseases, idition, a triple-dose preparation into	Every 10 days  Every 10 days  d vary, according to instructions, based on  t is essential to double the dosage of the fir				
How to use	Type of crop Floriculture Nurseries of young p Leafy greens The number of intervethe context, type of croposes can be diluted In the case of crops a 3 interventions. In additional context and some context are context.	Dose per hectare  1.5 liters  1.5 liters  2.5 liters  entions and dosage are indicative ar op and its duration. in any volume of water. ffected by resistant plant diseases, idition, a triple-dose preparation into	Every 10 days  Every 10 days  d vary, according to instructions, based on  t is essential to double the dosage of the fir				
How to use	Type of crop Floriculture Nurseries of young p Leafy greens The number of intervethe context, type of cr Doses can be diluted In the case of crops a 3 interventions. In additional evaluation	Dose per hectare  1.5 liters  1.5 liters  2.5 liters  2.5 liters  2.5 liters  2.5 liters  2.6 liters  3.7 liters  3.8 liters  4.8 liters  4.9 liters  4.9 liters  5. liters  6.	Every 10 days  Every 10 days  nd vary, according to instructions, based on  t is essential to double the dosage of the firevention should be carried out or according				
How to use	Type of crop Floriculture Nurseries of young p Leafy greens The number of intervethe context, type of cr Doses can be diluted In the case of crops a 3 interventions. In additional evaluation	Dose per hectare  1.5 liters  1.5 liters  2.5 liters  2.5 liters  2.5 liters  2.5 liters  2.6 liters  3.7 liters  3.8 liters  4.8 liters  4.9 liters  4.9 liters  5. liters  6.	Every 10 days  Every 10 days  nd vary, according to instructions, based on  t is essential to double the dosage of the firevention should be carried out or according				
How to use	Type of crop Floriculture Nurseries of young p Leafy greens The number of intervethe context, type of cr Doses can be diluted In the case of crops a 3 interventions. In additional evaluation	Dose per hectare  1.5 liters  1.5 liters  2.5 liters  2.5 liters  2.5 liters  2.5 liters  2.6 liters  3.7 liters  3.8 liters  4.8 liters  4.9 liters  4.9 liters  5. liters  6.	Every 10 days  Every 10 days  d vary, according to instructions, based on  t is essential to double the dosage of the fir				
How to use	Type of crop Floriculture Nurseries of young p Leafy greens The number of interve the context, type of cr Doses can be diluted In the case of crops a 3 interventions. In add to technical evaluation It is miscible with all of	Dose per hectare  1.5 liters  1.5 liters  2.5 liters  2.5 liters  2.5 liters  2.6 and its duration.  3.6 in any volume of water.  4.7 iffected by resistant plant diseases, it dition, a triple-dose preparation interest.  5. bother products and can be distributed.	Every 10 days  Every 10 days  and vary, according to instructions, based on  t is essential to double the dosage of the fire ervention should be carried out or according and either via root or foliar application.				
How to use	Type of crop Floriculture Nurseries of young p Leafy greens The number of interve the context, type of cr Doses can be diluted In the case of crops a 3 interventions. In add to technical evaluation It is miscible with all of	Dose per hectare  1.5 liters  1.5 liters  2.5 liters  2.5 liters  2.5 liters  2.6 and its duration.  3.6 in any volume of water.  4.7 iffected by resistant plant diseases, it dition, a triple-dose preparation interest.  5. bother products and can be distributed.	Every 10 days  Every 10 days  and vary, according to instructions, based on  t is essential to double the dosage of the fire ervention should be carried out or according and either via root or foliar application.				
How to use	Type of crop Floriculture Nurseries of young p Leafy greens The number of interve the context, type of cr Doses can be diluted In the case of crops a 3 interventions. In add to technical evaluation It is miscible with all of	Dose per hectare  1.5 liters  1.5 liters  2.5 liters  2.5 liters  2.5 liters  2.5 liters  2.6 liters  3.7 liters  3.8 liters  4.8 liters  4.9 liters  4.9 liters  5. liters  6.	Every 10 days  Every 10 days  and vary, according to instructions, based on  t is essential to double the dosage of the fire ervention should be carried out or according and either via root or foliar application.				
How to use	Type of crop Floriculture Nurseries of young p Leafy greens The number of interve the context, type of cr Doses can be diluted In the case of crops a 3 interventions. In add to technical evaluation It is miscible with all of Even in higher doses to	Dose per hectare  1.5 liters  1.5 liters  entions and dosage are indicative are op and its duration. in any volume of water. ffected by resistant plant diseases, in dition, a triple-dose preparation interests. bother products and can be distributed than indicated, it never damages the	Every 10 days  Every 10 days  Individual of the dosage of the first ervention should be carried out or according to either via root or foliar application.				
How to use	Type of crop Floriculture Nurseries of young p Leafy greens The number of interve the context, type of cr Doses can be diluted In the case of crops a 3 interventions. In add to technical evaluation It is miscible with all of Even in higher doses to Any changes in color a	Dose per hectare  1.5 liters  1.5 liters  entions and dosage are indicative are op and its duration. in any volume of water. ffected by resistant plant diseases, in dition, a triple-dose preparation interests. other products and can be distributed than indicated, it never damages the lare due to variation in the Earth's manages are due to variation in the Earth's manages.	Every 10 days  Every 10 days  Individual of the dosage of the first ervention should be carried out or according to either via root or foliar application.				
How to use	Type of crop Floriculture Nurseries of young p Leafy greens The number of interve the context, type of cr Doses can be diluted In the case of crops a 3 interventions. In add to technical evaluation It is miscible with all of Even in higher doses to Any changes in color a	Dose per hectare  1.5 liters  1.5 liters  entions and dosage are indicative are op and its duration. in any volume of water. ffected by resistant plant diseases, in dition, a triple-dose preparation interests. other products and can be distributed than indicated, it never damages the lare due to variation in the Earth's manages are due to variation in the Earth's manages.	Every 10 days  Every 10 days  d vary, according to instructions, based on  t is essential to double the dosage of the firervention should be carried out or according ed either via root or foliar application.				
	Type of crop Floriculture Nurseries of young p Leafy greens The number of interve the context, type of cr Doses can be diluted In the case of crops a 3 interventions. In add to technical evaluation It is miscible with all of Even in higher doses to	Dose per hectare  1.5 liters  1.5 liters  entions and dosage are indicative are op and its duration. in any volume of water. ffected by resistant plant diseases, in dition, a triple-dose preparation interests. other products and can be distributed than indicated, it never damages the lare due to variation in the Earth's manages are due to variation in the Earth's manages.	Every 10 days  Every 10 days  Individual of the dosage of the first ervention should be carried out or according to either via root or foliar application.				
How to use  Item Code	Type of crop Floriculture Nurseries of young p Leafy greens The number of interve the context, type of cr Doses can be diluted In the case of crops a 3 interventions. In add to technical evaluation It is miscible with all of Even in higher doses to Any changes in color a	Dose per hectare  1.5 liters  1.5 liters  entions and dosage are indicative are op and its duration. in any volume of water. ffected by resistant plant diseases, in dition, a triple-dose preparation interests. other products and can be distributed than indicated, it never damages the lare due to variation in the Earth's manages are due to variation in the Earth's manages.	Every 10 days  Every 10 days  d vary, according to instructions, based on  t is essential to double the dosage of the firervention should be carried out or according ed either via root or foliar application.				

**AXS M31** di Zambanini Silvana advanced bio-formulation for agriculture

M32FV-Mo/5L

Località Deggia, 28 38078 San Lorenzo Dorsino (TN) - I Tel. (+39) 0465 734591

5 liter tank

BioAksxter® M32 Photovoltaic Greenhouse

info@axsm31.com info@bioaksxter.com www.bioaksxter.com

5.150 kg